

UNIVERSITY of TASMANIA



Institute for Marine and Antarctic Studies

TASMANIAN SCALLOP FISHERY ASSESSMENT 2021

Jayson Semmens, Graeme Ewing, Ryan Day, Sarah-Jane Pyke, Samantha Twiname

March 2023



Institute for Marine and Antarctic Studies, University of Tasmania, Private Bag 49,
Hobart TAS 7001

Enquires should be directed to:

Dr Jayson Semmens

Institute for Marine and Antarctic Studies
University of Tasmania
Private Bag 49, Hobart, Tasmania 7001, Australia

Jayson.Semmens@utas.edu.au

Ph. (03) 6226 8275

The authors do not warrant that the information in this document is free from errors or omissions. The authors do not accept any form of liability, be it contractual, tortious, or otherwise, for the contents of this document or for any consequences arising from its use or any reliance placed upon it. The information, opinions and advice contained in this document may not relate, or be relevant, to a reader's particular circumstance. Opinions expressed by the authors are the individual opinions expressed by those persons and are not necessarily those of the Institute for Marine and Antarctic Studies (IMAS) or the University of Tasmania (UTAS).

© The Institute for Marine and Antarctic Studies, University of Tasmania 2023.

Copyright protects this publication. Except for purposes permitted by the Copyright Act, reproduction by whatever means is prohibited without the prior written permission of the Institute for Marine and Antarctic Studies.

Contents

Contents	1
Executive Summary	2
1. Introduction	4
History of the TSF	4
Key research	14
Overview of Management of the Tasmanian Scallop Fishery	19
Recreational fishery.....	22
2. Fishery peer review	23
3. 2021 Pre-season survey	25
Introduction and survey methods	25
Survey results by region	27
Biomass estimates	38
Conditions for opening the 2021 commercial scallop season.....	40
4. Economic data	41
5. Warm water event	42
6. Stock status	43
7. Bycatch and protected species interaction	44
8. Conclusions and recommendations	46
9. References	48

Executive Summary

STOCK STATUS	RECOVERING
The Tasmanian Scallop Fishery (TSF) is managed with a harvest strategy where surveys are undertaken to estimate abundance and decision rules are used to open areas to fishing with total allowable catches (TACs) based on the estimated abundance. Biomass in the Tasmanian Scallop Fishery (TSF) is historically overfished (Caton and McLoughlin, 2004), with recruitment and production levels now affected. In 2013, 2014 and 2015, surveys generally found low scallop densities and limited evidence of successful recent recruitment but did identify two beds (one on the north-west coast and the other on the east coast) containing commercial quantities (Ewing <i>et al.</i>, 2016). Surveys in 2016 and again in 2017 generally only found very low levels of scallop abundance and limited evidence of successful recruitment, with no area considered to contain commercially viable quantities in either year (Ewing <i>et al.</i>, 2017; Semmens <i>et al.</i>, 2018). This includes the east and north-west coast beds fished in 2013-2015, which appeared to have been fished down to a commercially unviable density, with no subsequent recruitment evident. Given the results of the 2016 and 2017 surveys, there was a low expectation that conducting pre-season surveys in 2018 and 2019 would yield the presence of commercially viable scallop beds, and as such, pre-season surveys were not conducted. In 2020 a pre-season survey was conducted, and the survey results were dominated by low to moderate densities of legal sized scallops, several areas demonstrated signs of recruitment that suggest the possibility of supporting a tightly managed commercial fishery in the near- to medium-term. The 2021 pre-season survey consisted of 147 targeted dredge tows and 48 video surveys. Survey results showed low to moderate densities of legal sized scallops, with patchy high densities and variable discard rates. Fishing mortality is managed with the aim of restricting catches to beds of mature scallops near the end of their lifespan. The combination of the harvest strategy and depleted biomass has led to a history of closures due to low abundance. In recent times, the fishery was closed between 2000-2002 and again between 2009 and 2010. Areas with commercial density of scallops towards the end of their lifespan were opened to fishing each year between 2013 and 2015. The fishery was closed between 2016 and 2020. The TSF was partly opened in 2021, the first season to open since 2015, with the TAC set at 1494.9 t or 145 kg/unit. However, only 44.5 t were landed from the open area due to a major die-off event post survey. The remainder of the state continued to be closed. On the basis that biomass has increased during the 2016-20 closure due to wide-scale recruitment events, and current restrictions are effectively limiting fishing mortality, the TSF is classified as recovering.	
STOCK	Tasmanian Scallop Fishery
INDICATORS	Size structure, density and catch

The TSF targets the commercial scallop (*Pecten fumatus*), one of three species naturally occurring in Tasmania. The SE Australian commercial scallop fishery began in Tasmania in the early 20th century and has exhibited a classic boom/bust trajectory with a combination of high recruitment variability and repeated overfishing and serial depletion leading to periods of very high effort and catches followed by extended closures for stock recovery. The TSF extends to 200 nm from the Tasmanian coast, with the exception of Bass Strait, where its jurisdiction covers 3-20 nautical miles offshore (DPIPWE, 2005). The fishery is managed by an individual transferable quota (ITQ) program and several input controls.

In comparison with historical levels, Tasmanian commercial scallop stocks are severely depleted and since 2003, the TSF has been regulated under a spatial management approach, referred to as a “paddock” fishery, whereby all areas are closed to fishing, except that certain areas can be opened after a stock assessment is completed. This approach is intended to promote stock re-building whilst allowing continuity of employment and supply for a small fishing fleet.

Sampling for the pre-season stock assessment has been conducted by selected scallop fishing vessels (issued with survey authorisations), and scallop stocks are assessed against criteria including:

- Population structure – particularly size where >80% of scallops landed must be greater than the minimum legal size (90 mm shell length (SL); animals of this size are approximately 3+ years of age and have had at least two major spawning’s).
- Commercial viability – including catch rates, market suitability, estimated costs of fishing and meat recovery (<85 scallop meats per kg).

The Scallop Fishery Advisory Committee (ScFAC) considers the pre-season-stock assessment and then provides advice to the ministry on which, if any, “paddocks” should be open for fishing. Members of ScFAC include representatives from the scallop fishing and processing industry, the research community, the relevant federal and state fisheries management agencies, and the marine conservation sector; and reflect the co-management framework for the management of the Tasmanian scallop fishery.

Sampling for the 2016 and 2017 pre-season stock assessments (Ewing *et al.*, 2017; Semmens *et al.*, 2018) on the east, north east and northwest coasts of Tasmania indicated low densities of scallops in all regions. Further, these surveys found limited evidence of successful recruitment. These findings were also supported by an IMAS fishery-independent video survey on the east coast in 2017. Based on these previous pre-season surveys, ScFAC advised the minister not to open the Tasmanian commercial scallop fishery in both 2018 and 2019 and stock status of the TSF remained classified as depleted. However, the pre-season dredge survey conducted in May-June 2020 (Ewing and Semmens 2020) offered evidence of recruitment across several areas of the fishery. While this has raised the possibility of stock recovery in these areas, based on the overall low to moderate densities of legal sized scallops in the dredge survey and the results of an IMAS fishery-independent video survey conducted in Great Oyster Bay (Semmens *et al.*, 2020), ScFAC advised the minister not to open the Tasmanian commercial scallop fishery in 2020, but to further monitor recovery prior to the 2021 season.

The 2021 survey subsequently covered three Flinders Island beds (Scallop Fishery Area 3) with an estimated legal-sized (> 90 mm SL) biomass of 17,145 t, one central east coast bed (Scallop Fishery Area 4) with an estimated legal-sized biomass of 384 t and one lower east coast bed (Scallop Area 5) with an estimated legal-sized biomass of 9,366 t (Ewing *et al.* 2021). Overall, the 2021 pre-season survey showed low to moderate densities of legal sized scallops, with patchy high densities and variable discard rates and a partial opening was recommended by ScFAC in 2021, the first season to open since 2015.

Scallop Area 3 opened to fishing in September 2021, the first season since 2015, with the TAC set at 1494.9 t. However, there had been a major die-off post survey, most likely due to a warm-water incursion in the Flinders Island area and only 44.5 t was landed due to poor quality and recovery. The remainder of the Scallop Fishery Areas remained closed.

1. Introduction

History of the TSF

The Tasmanian scallop fishery began as a recreational fishery in the Derwent Estuary in the early 1900s. A commercial fishery developed in 1919 and in the ensuing 5 years expanded rapidly and moved to target beds in the D'Entrecasteaux Channel. At the time the loss of stocks in the Derwent were attributed to flooding and predation by crabs and starfish, however, over-fishing, industrial pollution and siltation from land-use practises are likely to have played a role in subsequent declines.

The northern beds in the D'Entrecasteaux Channel were closed amid concerns of over-fishing in 1925, then were re-opened, then were closed again in 1930, and again re-opened. Sharp declines in catches in the Channel in the 1940s were attributed to over-fishing and poor recruitment, and the fishery began exploiting east coast stocks (Figure 1). By the 1970s east coast stocks were severely depleted and effort shifted to large scallop beds off the Furneaux Island Group in Bass Strait by the late 1970s (Figure 1). Landings and numbers of vessels increased rapidly (Figures 1 & 2), with 12,000 t (live weight) taken in Bass Strait in 1983 by 231 vessels. By the late 1980s these beds were also severely depleted; in fact there were virtually no productive scallop grounds left in southern Australian (Caton and McLoughlin, 2004) (Figures 1 & 2).




Figure 1. Commercial catches, as total meat weights, of *P. fumatus* from Bass Strait and Tasmanian and Victorian waters from 1928 to 1989. The years when new beds were first exploited are indicated by arrows (Young *et al.*, 1988).




Figure 2. Commercial scallop catch (shell wt.) within the Tasmanian scallop fishery.

Stricter management arrangements commenced in the 1980s including catch history for access, allocation of fishing units, limits on vessel numbers, and fishery closures. The fishery was totally closed for 8 years from the end of 1987 until 1995 (Figure 2) to promote the rebuilding of the scallop stock. Partial recovery of the stock in some areas led the fishery to be opened for short seasons in 1995, 1996, 1998 and 1999, although the fishery again closed between 2000 and 2002 (Figure 2).

Following the introduction of ‘bag’ quotas in the 1990s, transferable units were introduced in 2000 to encourage restructuring in the fishery. A total of 10,730 scallop units were issued to operators and these remain in place as a fundamental component of the fishery. To limit the level of catch increasing as a result of activating effort from latent units and licences, the quota unit value was reduced effectively reducing the ‘bag’ quota unit value based on volume (equating to around 950 kg) to a weight-based value “kilogram scallop unit” of 500 kg. These measures reduced the total potential catch from 10,400 t (if all units were activated and used) to a more conservative level of 5,350 t.

In 2002 extensive areas of scallops were identified off Flinders Island (Furneaux Island group), however, the fishery remained closed as most of these scallops were undersize. Regardless, this generated a high level of interest in the fishery indicating that it was highly probable that most scallop licences would be activated, and most scallop units would be seasonally transferred. This created the potential for catch to exceed 4200 t and the trigger point relating to capacity of active licences being reached. DPIPWE took a proactive approach, and the scallop unit value was reduced from 500 kg to 400 kg prior to the commencement of the fishery opening in June 2003. The change reduced the maximum potential catch from 5,350 t to 4,280 t. While the industry accepted that there were no guarantees that they would catch their quota each season, DPIPWE considered it created an expectation for the opening of all areas that fulfilled the discard criteria in the immediate season.

After the fishery experienced considerable difficulties in re-establishing markets after being a long period of total fishery closure (1999 to June 2003), a spatial management strategy was refined to focus on providing for continuity of fishing seasons. This continues to be a primary aim of the management of the fishery. The intention was to develop harvesting plans for potential open areas, ranking the order in which particular areas could be opened including, where feasible, reserve areas to ensure fishing seasons 2-3 years into the future.

In 2003, three areas were opened under the spatial management strategy, east of Flinders Island, Banks Strait (stretch of water between north-eastern Tasmanian and the Furneaux Island group) and Eddystone Point (east coast), with the 30 participating operators catching 3324.5 t of the 4146.4 t TAC (10366 units at 400kg per unit). Similarly, in 2004, three areas were opened, St Helens Point to

Schouten Island (east coast), then Eddystone Point and Marion Bay (east coast), with all of the 4146.4 t TAC caught by the 24 participating operators (see Table 1). For both seasons, period and trip limits were used to limit catch.

Table 1: Summary of the Tasmanian scallop fishery 2003-2015. *Denotes that operators were also able to apply for an extra 50kg/unit to be caught off Flinders Island in 2005.

Year	TAC	Operators	Days fished	Catch tonnes
2003	4146.4	30	1010	3324.5
2004	4146.4	24	1402	4159.5
2005	4146.4*	25	1642	4329.0
2006	4146.4	24	1386	4359.3
2007	4253.2	20	634	1390.9
2008	4253.2	15	480	489.8
2011	2552.0	10	102	103.3
2012	1163.0	13	373	790.2
2013	1489.0	14	552	1226.0
2014	1240.0	6	196	489.0
2015	1033.0	11	261	781.2
2021	1494.9	2	8	44.5

In 2005, a new management plan was introduced which allowed the Minister to set and alter the quota unit value by public notice. The alteration in unit value effectively sets the TAC for any particular year and is based on the best available stock information at that time. DPIPWE intended to use this management tool in combination with the harvesting plans discussed above to maximise the likelihood of continuous fishing years. In addition, industry restructuring resulted in a reduction of the number of licences to 92 by 2005. In 2005, the Tasmanian Aquaculture and Fisheries Institute (TAFI) advised that surveys had not indicated signs of significant recruitment of scallops and that the amount of known harvestable scallop stocks was likely to be less than the cumulative TAC over the next few years. To assist in meeting the objective of continuous fishing years, and to continue stock rebuilding, DPIPWE adopted a precautionary approach and reduced the TAC to 3628.1 t (by altering the unit value to 350 kg, with 10366 units) for the 2005 season.

Four areas were opened in 2005, Eddystone Point, Marion Bay, Paddy's Head to Schouten Island (east coast), and east of Flinders Island. Surveys conducted before fishing demonstrated that the scallop abundance east of Flinders Island was declining rapidly through natural mortality. In response, DPIPWE allocated an additional 50kg to the unit values, taking it back up to 400kg per unit, with operators also able to apply for an extra 50kg/unit to be caught off Flinders Island. A temporary legal minimum size of 80 mm shell length was also set for the east coast of Flinders Island. These measures were intended to ensure that fishers could maximise the east Flinders Island beds before they were completely lost through natural mortality. For all areas combined, 4329.0 t was caught in 2005, by 25 operators.

In 2006, White Rock (east coast) was initially opened, then Cape Lodi to Wardlaws Point (east coast). The 24 participating operators caught all the 4146.4 t TAC. Figures 3 to 6 below show the catch and effort for 2006 compared to more recent seasons (2011-2015), as 2006 was the first year that White Rock was fished since the reopening of the fishery in 2003. White Rock has been the most consistent site of the fishery in recent years (2011-2015) (Figure 3) but has produced significantly smaller catches than those in 2006. In 2007, initially a small area around Eddystone Point was open, with a larger area in Banks Strait subsequently opened. Only 1390.9 t of the 4253.2 t TAC was caught, marking the beginning of the first downturn in catches, corresponding to declining stocks, since the fishery re-opened in 2003. In 2008, the stocks declined further, and there was no significant recruitment. However, two areas were opened, one-off Bicheno (east coast) and another in Marion Bay. Pre-season scallop surveys found the two open areas contained low densities of large scallops

in good condition. The two open areas had been fished a number of times since 2003 and the scallops available for harvest were residual scallops from previous seasons with no evidence of recruitment to these beds. Despite this, the fishery was opened with a TAC of 4253.2 TAC, but a ministerial warning was issued that it was highly unlikely the TAC would be taken. This warning proved correct, with only 489.8 t caught by the 15 participating operators.

The Tasmanian fishery was closed to fishing in 2009–10 due to low abundance and small average sizes (below the minimum size).

The fishery opened again in 2011 (see Figures 3-6) after a survey in White Rock indicated some stock recovery in this region of the fishery. The TAC was set at 2552 t, however, after opening the fishery, it was found that the beds had suffered a large die-off, and very little fishing took place in the White Rock region (Figure 3). Only 103 t catch was taken in 2011, and this came from a state-wide survey that failed to identify significant new scallop beds.

In 2012 Marion Bay was initially opened with a TAC of 525 t. The White Rock area was opened later, and the TAC increased to 1163 t, however, the season was abandoned due to a toxic algal bloom on the east coast, with a total of 790 t of scallops landed for the season by the 13 participating operators.

In 2013, 2014 and 2015 (see Figures 3-6) industry surveys found low scallop densities and limited evidence of successful recent recruitment other than two beds with commercial quantities. One was in the Circular Head region on the north-west coast and the other in the White Rock region. These areas were subsequently opened to fishing each year between 2013 and 2015 (Figure 3). A lower minimum size limit (85 mm SL) was applied to the north-west bed given the historically slow growth rates in this region and the fact that most scallops of this size are still 3+ years old (Martin *et al.*, 1988).




Figure 3. Distribution of catch (tonnes) for 2006 and for the 2011 to 2015 seasons by block.




Figure 4. Distribution of effort (hours) for 2006 and for the 2011 to 2015 seasons by block.




Figure 5. Regional CPUE (kg/hr) for 2006 and for the 2011 to 2015 seasons by block.




Figure 6. Catch (tonnes), Effort (hours), and mean CPUE (kg/hr) for 2006 and for the 2011 to 2015 seasons. CPUE error bars are standard error.

A voluntary industry closure was implemented in July 2013 for a portion of the east coast bed when a large aggregation of sub-legal scallops was discovered in the northern section of the fishing area. The 2013 season was originally opened with a TAC of 638 t, which was later increased to 1063 t, and then to 1489 t, following in-season surveys, with 1226 t harvested by the 14 operators participating. Similarly, a TAC of 620 t was set in 2014, but later increased to 1240 t. However, only 489 t was caught, with 6 operators participating. Most of the catch (404 t) came from the Circular Head region, which had average scallop densities of almost twice those of the White Rock region when surveyed. Again in 2015, the initial TAC (620 t) was increased during the season, with only 781 t of the final 1033 t TAC taken, with 11 operators participating. Seven hundred and seventeen t came from the White Rock region, which had average scallop densities of almost twice those of the north-west bed when surveyed. However, it should be noted that two different vessels returned different scallop densities and size frequency distributions for the same area of the White Rock region surveyed in 2015, with the second survey showing a lower density and greater percentage of sub-legal scallops (greater than the 20% allowed under the decision rules).

Through 2014–17 market supply was maintained from the Commonwealth Bass Strait central zone scallop fishery (BSCZSF) which had significantly greater stocks than those for the Tasmanian fishery at this time. Most licensees in the Tasmanian fishery also operate in the Commonwealth fishery. As such, some of the uncaught TAC in the Tasmanian fishery during this period may have been caused by a shift in effort to Commonwealth waters, where scallops were more abundant. Given the more abundant resource in the BSCZSF, industry state-wide surveys were only conducted within known scallop beds on the state's east coast in 2016 (White Rock and Marion Bay regions), with no knowledge of the status of beds elsewhere in the state. In 2017, the White Rock and Marion Bay regions were again surveyed, along with Circular Head and the Flinders Island regions. For both years,

the surveys generally only found very low levels of scallop abundance and limited evidence of successful recruitment, with no area considered to contain commercially viable quantities. This included the White Rock and Circular Head beds fished between 2013 and 2015, which appeared to have been fished down to commercially unviable densities, with no subsequent recruitment evident.

Based on these previous pre-season surveys, ScFAC advised the minister not to open the Tasmanian commercial scallop fishery in both 2018 and 2019 and stock status of the TSF remained classified as depleted. However, the pre-season dredge survey conducted in May-June 2020 (Ewing and Semmens 2020) offered evidence of recruitment across several areas of the fishery, including White Rock, the east coast of Tasmania and the Flinders Island region.

The 2020 dredge survey covered all six state-wide survey areas (note that sub-areas were removed post 2020, see Figure 7). Overall, low densities, suggested that Area 1 could not support a commercial dredge fishery at the time of the survey and that the area had not yet recovered to levels prior to it being fished between 2013 and 2015 (Circular Head Region). Densities in Area 2, the western region of Area 3, Area 5 and sub-area 6C yielded uniformly very low densities of legal-size scallops, with very few scallops encountered across all dredge tows. The average density of scallops in Area 4 was moderate, with a high discard rate at 90mm of 32%. Even though a few small beds of higher density were encountered inshore off Ansons Bay and the Gardens, the eastern side of Flinders Island (Babel Island and Pot Boil) yielded more consistent beds of a moderate density. While Area 4 yielded a much higher density of legal-sized scallops than when it was last sampled in the 2017 pre-season survey, the 2020 distribution of scallops was patchy with a strong representation of undersize classes. This evidence of new recruits across Area 4 was a potential good sign for the future of the fishery.

White Rock (sub-area 6A) survey tows yielded an average density of 21 kg/1000 m², with a discard rate of 7.5%. Densities in the Marion Bay (6B) sub-area were lower than White Rock (6A), with an average density of 8.3 kg/1000 m². The discard rate of 15.6% in sub-area 6B was twice that of 6A. The improved density of scallops encountered in 6A and 6B in the 2020 survey, compared with the last two pre-season surveys, is likely to be due to four years of stock-rebuilding from the closure of the fishery since the 2015 season. However, despite the increase in density since the 2017 survey, the moderate (6A) and low (6B) density areas were highly spatially restricted, i.e., patchy, as was likely the limited recruitment in these areas. Along with the dredge survey, a towed underwater video camera was used to assess scallop densities and size frequency in survey sub-area 6A (White Rock) in May and June 2020 (Semmens *et al.* 2020). Higher density sites were highly spatially restricted, i.e., patchy. Size structure from video transects showed strong representation of sub-legal scallops (58.9% at the 90 mm legal-size) and a mean size <90 mm. Along with the discreet patches of higher densities, the high proportion of sub-legal scallops throughout the survey area, clearly demonstrated that there has been recruitment during the closure of survey sub-area 6A (and the entire TSF) since the 2015 fishing season. Furthermore, there are potential good signs for the future, with clear evidence of new recruits.

While the 2020 surveys raised the possibility of stock recovery around the state, based on the overall low to moderate densities of legal sized scallops in the dredge survey and the results of the IMAS fishery-independent video survey conducted in Great Oyster Bay, ScFAC advised the minister not to open the Tasmanian commercial scallop fishery in 2020. It should be noted that since 2014 (including 2020) market supply has been maintained by the Commonwealth Bass Strait central zone scallop fishery (BSCZSF), which has held significantly greater stocks than those for the Tasmanian fishery during this time period. The Victorian Ocean Scallop Fishery was also open in 2020.

The 2021 dredge survey covered several locations of the east coast of Tasmania, specifically White Rock, The Gardens, Pot Boil and North and South Babel. Only targeted surveys were conducted in 2021, no exploratory dredge surveys were conducted.

White Rock targeted dredge survey tows yielded an average density of legal-sized scallops of 15.6 kg/1000 m² (median 14.2 kg/1000 m²), with a discard rate of 5.67% (Ewing *et al.*, 2021). Results from the video survey yielded an average density of legal-sized scallops of 30.4 kg/1000 m² (median 27.2 kg/1000 m²), with a discard rate of 30.5% (Semmens *et al.*, 2023). Higher representation of smaller scallops in the video survey is expected and reflects the size selectivity of commercial scallop dredges.

The low to moderate densities of legal-sized scallops in the 2021 survey were similar to those encountered in the 2020 pre-season survey. The lower proportions of sub-legal scallops and higher average sizes relative to the 2020 survey reflect growth into the fishery of the standing crop of scallops but may also indicate reduced stock recruitment.

A low dredge survey discard rate, a decline in the video survey discard rate and high average size indicate that the White Rock region has significantly less undersize scallops in 2021. However, given the lack of recent recruitment, low to medium densities of legal-sized scallops and the history of openings and closures in this area, the merits of opening White Rock in 2021 needed to be given careful consideration, noting the depleted status of the Tasmanian scallop fishery. As such, ScFAC advised the minister not to open the White Rock region (Scallop Area 5) and instead Scallop Area 3 (eastern side of Flinders Island) opened to fishing in September 2021, the first season since 2015, with the TAC set at 1494.9 t. However, there had been a major die-off post survey, most likely due to a warm-water incursion in the Flinders Island area and only 44.5 t was landed due to poor quality and recovery. The remainder of the Scallop Fishery Areas remained closed.

Key research

Over the years, several large research projects have been undertaken by IMAS to facilitate management of the TSF.

FRDC 2012/027: Determining when and where to fish: Linking scallop spawning, settlement, size and condition to collaborative spatial harvest and industry in-season management strategies.

This project defined timing of scallop spawning based on gonad condition, and hence potential settlement of recruits across the different populations/beds of the fishery and determine any differences in spawning potential among scallop beds/locations. Additionally, this project defined differences in spawning potential between scallops ranging from 80 to 90 mm SL and assessed the size limits used to define a bed as commercially viable across the three southeast Australian jurisdictions. Furthering our understanding of growth rate in several fishing locations across all three jurisdictions allowed better management of individual beds and use of scallop condition in setting season openings and closings (Semmens *et al.*, 2019).

Changes in muscle weight, gonad weight and meat recovery weight (combined meat and gonad weight) varied through the season and the gametogenic cycle, but this relationship is affected by year. There was no common trend in changes in muscle, gonad and combined weight across areas, instead the changes are area-specific and year-specific and can vary considerably. The differences were of such a magnitude that each area requires research prior to harvesting. This is particularly relevant in the TSF, when only relatively small areas of the fishery are opened at any one time and the remainder of the fishery is closed, although the adaptive in-season management model in this fishery (i.e., open area boundaries can be changed during the season) can overcome this issue in some circumstances. In the BSCZSF and OSF, given only relatively small areas of the fishery are closed during the fishing season or the entire fishery is opened, respectively, there may be a greater opportunity for scallop fishers to find beds with higher muscle, gonad and combined weights, although this of course relies on fisher behaviour and is dependent on the number of different beds available, with limited to no scallop beds available for harvest in the OSF for over two decades. Furthermore, without adaptive in-season management, which is the case in the BSCZSF (i.e., spatial closures are not adjusted during the fishing season), there is the potential for the best quality scallops to be 'locked-up' in spatial closures for the entire season.

Maturity stages identified macroscopically did not consistently match the maturity stages identified by histological sampling. Apart from macroscopic stage 2, which comprised scallops with predominantly gonads in the developing stage, the other macroscopic stages showed a mixture of reproductive stages. Therefore, while the macroscopic staging scheme is useful to derive a general indication of gonad condition, it does not accurately reflect the maturity stage in the ovary.

Based on microscopic observations compared to macroscopic examination of gonads, three visual stages are described based on the morphological appearance of the gonad to the naked eye: Developing or spent; maturing or atretic (reabsorbing eggs as spawning is delayed); and partially spawned. Fishing in the Commonwealth predominately takes place in the beds or regions surveyed before the season commences. Similarly, the TSF only opens to fishing surveyed areas that meet the management plan. While the OSF does not use surveys to determine open areas, as it is not a spatially managed fishery, fishing generally occurs in traditional areas, which will have variable gonad development and spawning timings within and between them. As such, the simple three stage visual classification system developed in this project is useful to both scallop resource managers and industry, as part of in-season management strategies, to define the overall reproductive stage of scallops and predict timing of spawning, thus assisting in the best condition scallop beds being fished sequentially throughout the season.

Collection of data on scallop condition, reproductive stages and settlement rates collectively can help inform best timing for season opening and closing dates in each location. The information available from this study and previous studies, suggest that the Lakes Entrance region, which comprises the majority of the OSF, would profit from an early start during winter. However, note that the OSF is currently considered depleted and has not had significant catches in over two decades. This in part

may be attributable to the fact that the fishery has historically been open continuously throughout the year, including the settlement period. At White Rock in the TSF, starting the fishery in September would appear more beneficial in terms of harvesting the best product, although this may not fit best with protecting newly settled scallops, and may in part explain why this area has not supported a fishery in recent years and is now classified as depleted. At the Bass Strait site in the eastern section of the BSCZSF, the best time to fish appears to be spring and summer/autumn. Fishing up to the closing date of December 31 may not fit best with protecting newly settled scallops, with the major settlement period occurring in spring, and again may in part explain why this area has not been viable in recent years. At King Island, in the BSCZSF, the best time to fish appears to be spring and summer, however, settlement occurs from approximately November to January.

Fecundity increased exponentially with SL and modelling predicted that a scallop measuring 90 mm in shell length would be 13 and 25% more fecund than an 85 and 80 mm scallop, respectively. Furthermore, an 80 mm scallop would be 44% more fecund than a scallop measuring 70 mm in SL. Scallops measuring 100 mm in SL would produce 32% more eggs than a scallop measuring 90 mm. These differences are less dramatic than previous findings where 3+ years old scallops measuring ~90 mm SL shed (3.5 million eggs on average) compared to 2 million eggs shed by scallop measuring ~83 mm SL (a 57% difference compared to 19% estimated in this study). This result of the current study showing a much smaller difference in fecundity in scallops of various sizes compared to previous findings, is a very important finding in relation to the decision rules around scallop harvest, particularly the under-sized discard rate rule and the two spawnings criteria which states that scallops should be allowed a minimum of two major spawning events before being harvested. Scallops that are 85-95 mm SL are 3+ years old and have had two major spawning and thus contributed significantly to potential recruitment. However, given the relationship between fecundity and SL demonstrated in this study, which shows a 3-fold decline in the difference between fecundity of an 83 and 90 mm SL scallop compared to the previous research, the size limits are very conservative. As such, the use of 85 mm SL still allows the scallops to have produced two major spawnings before harvest, with relatively little difference between the fecundity of 85 and 90 mm SL scallops (13%). However, it should be noted that in regions that have very low biomass or are recovering from being depleted (e.g., TSF), this additional 13% could be significant, and a highly conservative approach may be warranted. Furthermore, it should be noted that the 80 mm SL size limit used for the decision rules in the OSF is likely not appropriate, as it is outside of the size range for the two major spawnings criteria and should be revisited, with this low size limit perhaps contributing to the long history of limited biomass and recruitment in the fishery.

Differences in shell morphology were evident among regions, with significant differences between the standardised height for a standard scallop measuring 90 mm SL. However, differences in morphology were more evident among locations when comparing shell widths for standard scallops measuring 90 mm SL. Northwest Tasmania (TSF) and Great Bay (TRSF) had comparatively thinner individuals, followed by King Island (BSCZSF), Banks Strait (TSF), Marion Bay (TSF) and White Rock (TSF). Scallops from Babel Island (TSF and BSCZSF) showed no significant differences in shell width with Eddystone (TSF) or the Bass Strait (BSCZSF) site. Scallops located in Victoria (OSF) had the thickest scallops. For fisheries management purposes it is interesting to determine if scallops with greater SW also have greater muscle and/or gonad weights. Indeed, the Victorian (OSF) site had the thickest (deepest) scallops, and these scallops generally had the heaviest muscles, gonads and combined weights in winter of all the regions. Additionally, other thick scallop regions, Isthmus Bay (TRSF) and Eddystone Point (TSF), also had heavy muscles, gonads and combined weights in winter. Furthermore, the relatively thick scallops from the Bass Strait (BSCZSF) site had heavy muscles, gonads and combined weights in autumn. Fishing these areas (when opened) in the seasons noted could increase commercial yields.

Scallops at different sites showed variable mean growth increments depending on initial mean size of cohorts. There was no obvious growth pattern on the latitudinal gradient. For instance, sites at the extreme north (North Flinders and King Island, both in the BSCZSF) and south (Great Bay, TRSF) of Tasmania showed average mean growth increments. Low and high values of growth were observed in sites that are close to each other in the BSCZSF (1.9 and 9.20 mm/year for King Island Middle and King Island 2 respectively). Therefore, growth variations seem to be associated with local factors rather than factors linked with large spatial scale change, which has also been observed for other species of scallop. This growth analysis has shown that there is great variation in growth rates of

commercial scallop across the traditional fishing areas within the southeast of Australia, with great variation even prevalent between beds in the same area, e.g., King Island (BSCZSF) and East Flinders Island (TSF).

Importantly, however, this analysis has shown that the fishing areas examined can be generally grouped into three general groups: rapid growers; moderate growers; and slow growers. Rapid growers will be younger than their shell length indicates, so those scallops may not be 3+ at 85-95 mm SL and may not have had three major spawnings. As such, despite the relationship between size and fecundity showing that the 90 mm size limit is very conservative, it may not be for these rapidly growing scallops, and perhaps a more conservative approach is needed, particularly as the size limit used in the BSCZSF, where the two North Flinders fishing areas are located, is 85 mm SL. Alternatively, if a validated aging technique can be developed for commercial scallops, this should be adopted to ensure scallops are only fished from the 3+ age class onwards. It is interesting to note that three rapid growing areas are North and Northwest Flinders (BSCZSF) and White Rock (TSF), all areas that have shown large reductions in biomass, with little or no recruitment in recent years. The TSF had plans to use an 85 mm SL size limit from 2020 onwards, however, based on this research, the 90 mm SL size limit was maintained.

Note that as previously mentioned, there is great variability with areas, with Flinders Island 1 (BSCZSF), which like North and Northwest Flinders is also situated north of Flinders Island, showing slow growth. Two other slow growing areas are at King Island (KI 2 and KI New, BSCZSF), with King Island Mid (BSCZSF) showing moderate growth. The slow growing scallops will be older than their shell length indicates, and as such 90 mm SL minimum size is likely to be very conservative. The King Island sites are in the BSCZSF, and as such currently managed under an 85 mm SL size limit, which would appear very appropriate, but likely to be highly conservative. This may be a key factor in the fact that this region has been maintaining very high biomasses despite the fishery operating in the region since 2014 and ~12500 t coming out of the area (west of 147 degrees east) in that time. Although note that the Northwest region in Tasmania is fished with an 85 mm SL size limit rule, as fishers nominated this area as a slow growing scallop area, however it has undergone a large decline in biomass, following no recruitment in recent seasons.

In those areas with slow growing scallops, closing beds based on the 20% discard rule may mean that some beds that have 80% or greater of the scallops within it having reached 3+ and having had at least two major spawnings may be inadvertently closed, and as such this rule will be very conservative in these areas. Conversely, the opposite will apply in fast growing areas, with beds that have less than 80% of the scallops within it having reached 3+ and having had at least two major spawnings being inadvertently opened, and as such this rule is not met in these, which could have an impact on the sustainability of fisheries in these areas. As such, defined use of the minimum size limit and 20% discard rule is not appropriate, and instead they should be used in conjunction with the known attributes of the beds within region to be fished and applied in an informed and sensible manner such that recruitment potential is not impacted. If a validated aging method can be developed for commercial scallops, the 20% discard rule will be able to be applied with greater confidence.

FRDC 2008/022: Establishing fine-scale industry based spatial management and harvest strategies for the commercial scallop fishery in South East Australia.

FRDC 2008/022 examined scallop stock structure, spawning biomass density/recruitment relationships, and the impacts of intensive fine spatial scale fishing on scallop communities with the aim of refining detailed spatial management/industry fine-scale management harvest strategies, such that they promote recruitment and minimise impacts on the broader environment (Semmens *et al.*, 2015).

This project demonstrated that the south east Australian commercial scallop population is largely a genetically homogeneous single population. However, there was some evidence of population structure within Bass Strait, which has implications for management of apparently genetically linked populations in separate management jurisdictions (Tasmania, Victoria, and Commonwealth). Evidence from this study suggested that these differences may be due the effect of ocean gyres that exist in the Bass Strait and may force self-recruitment of certain beds and genetic separation from the general population of the scallops in the Strait. It is important to note that these differences may be

due to genetic drift rather than spatial barriers to cross bed recruitment. This project also demonstrated that overall genetic exchange appears to be limited when distances exceed 300km and the finest scale at which genetic subdivision was found was approximately 100km. Importantly, this study demonstrates that appropriate scales of management should consider both long established patterns of dispersal and recruitment as indicated by population genetic structure, as well as short term patterns due to demographic heterogeneity. The genetic evidence indicates that stock structuring can occur within 100 km implying that yearly stock- recruitment dynamics are likely to exist on even smaller spatial scales. In other words, recovery of depleted scallop beds in the short term will be heavily influenced by recruitment from adjacent scallop beds rather than from distant beds. This study reinforces existing knowledge based on the modelling of currents in Bass Strait (Hammond *et al.*, 1994) indicating that cross seeding of scallop beds is rare. The scale of the current spatial management harvest strategies employed in both the Commonwealth and Tasmania would seem appropriate given the genetic evidence presented here. However, the harvest strategy employed in the Commonwealth, where a proportion of known beds cannot be fished, should perhaps be considered in Tasmania, given that localised recruitment appears to be the driving process.

In accordance with research in other benthic molluscs, this project found a strong indication that the density of recruits is related to the density of adults in the previous year. In the areas studied, recruit density increased by between 2 and 10 times for every single unit of adult density prior to spawning. The density of adult spawners also has an impact on the level of synchronicity between spawning adults. This study showed a difference in spawning intensity and synchronisation between sites of high and low densities and suggests that maintaining dense areas of adult scallops may increase the probability of recruitment, through increased spawning intensity. To this end, the protection of dense scallop beds has been incorporated into the Commonwealth Harvest Strategy, with at least part of a high density bed(s) found during surveys closed to fishing during the season. Again, this may well be a strategy worth adopting in Tasmania to promote recruitment.

Analysis of data from the Before-After-Control-Impact (BACI) study component of this project, which aimed to determine the impacts of commercial dredge fishing effort on the benthic communities found within scallop habitat, showed no significant effect on abundance or species composition whether or not scallop beds were fished, the level of fishing effort, the sampling year or the region sampled. This suggests that intensive fine spatial scale fishing associated with spatial management has no obvious short- to medium-term detrimental effects on scallop communities within the fishing grounds of the Commonwealth Bass Strait Central Zone Scallop Fishery (BSCZSF). As such, allowing for beds to potentially be rotationally fished after relatively short temporal closures, if they meet the harvest strategy decision rules, may have relatively little impact on the scallop communities.

However, it is important to note that the BSCZSF has been fished since the 1970s, and the small differences in species assemblages selective to dredge fishing between fished and non-fished sites found in this study may be due to 'historical impacts' the dredge fishery may have had on the benthic community. Repeated dredging over many years may have shifted the entire community to one which is more resilient and which can withstand dredge fishing pressure as has been found in other studies (Bradshaw *et al.*, 2001). Those species that are most affected by dredging may now be too rare to be effectively sampled using dredge surveys.

Furthermore, previous studies have also shown that changes in community structure following seasonal weather events can be more significant than those changes associated with fishing (Currie and Parry, 1996). Communities within the BSCZSF are more exposed to environmental variability than other regions where commercial scallops are or have been fished, perhaps further explaining why changes in species abundance were not observed in this study, but were observed for commercial scallops in inshore waters (see Currie and Parry, 1996). Despite this, heavily fished inshore regions also show resilience to fishing, with species abundances similar to those prior to dredging within six to nine months after fishing in the Port Phillip Bay Scallop fishery (Currie and Parry, 1996), which was fished close to continually for around 30 years.

FRDC 2005/027: Facilitating industry self-management for spatially managed stocks: a scallop case study.

FRDC 2005/027 established the capacity for industry to organise and implement surveys at both the scale of the fishery, and the scale of individual scallop beds (Harrington *et al.*, 2008a). These surveys are independent of direct research and management involvement. The population structure and abundance data that industry obtains during such surveys can be used by management to meet decision rules allowing the successful implementation of detailed spatial management strategies within the fishery. Furthermore, the development and use of electronic measuring and recording devices both simplifies and adds a level of credibility to the process of data collection, storage and analysis. The inclusion of industry in the data collection process of management also creates a sense of Industry ownership. In general, this improves the relationship and communications between all stakeholders in the fishery (industry, managers and research), and creates both an acceptance and level of understanding of the biological and economic benefits of detailed spatial management. This belief in the benefits of spatial management has directly led to industry empowerment, with much greater roles and responsibilities in the management of their fishery.

FRDC 2003/017: Juvenile scallop trashing rates and bed dynamics: testing the management rules for scallops in Bass Strait.

FRDC 2003/017 concluded that spatial closures in the management of commercial scallop stocks, where the majority of the fishery is closed to fishing and only small discrete regions of the fishery are opened to harvesting, offers a real prospect for providing continuity and sustainability for the fishery (Haddon *et al.*, 2006), especially when compared to conventional management. It also identified the very extensive data/stock information requirements of closed area spatial management (Haddon *et al.*, 2006; Harrington *et al.*, 2007). Without credible up-to-date stock information, scallop beds cannot be opened to harvesting, within season contingency plans cannot be formulated, and longer-term harvest strategies cannot be developed. This type of longer-term information and planning is essential in creating a level of certainty within the catching sector (industry), which in turn allows the development of processing infrastructure and domestic and export markets.

Overview of Management of the Tasmanian Scallop Fishery

The Tasmanian commercial scallop fishery extends to 200 nm, except Bass Strait, where jurisdiction covers 3-20 nautical miles offshore (DPIPWE, 2005) (see Table 1 for a summary of the fishery). The fishery is managed by an individual transferable quota (ITQ) program, which was implemented in 1986, as well as several input controls. Since 2003, the fishery has been managed under a spatial management approach, referred to as a paddock fishery, whereby all areas are closed to fishing, except that certain areas can be opened after a stock assessment is completed. The Fisheries (Scallop) Rules 2020 (current January 2021) is a collection of detailed fishery rules governing the fishery. A draft document, A Management of the Tasmanian Scallop Fishery – Policy and Decision Making Guidelines, describes the framework for management of this fishery, including goals, objectives, and strategies. This strategy specifies some main criteria that need to be satisfied for a specific area (paddock) to be opened (and remain opened) to fishing:

1. Minimum size or age – At least 80% of scallops must be greater than minimum legal size (90 mm SL) or at least 80% of scallops are ages 3+, corresponding to at least two major spawning opportunities. Alternative minimum size limits can be established, in cases where it can be established that scallops have had the opportunity to spawn at least twice prior to harvest.
2. Commercial viability – Potential fishing areas are prioritized based on anticipated economic returns, catch rates, market suitability, and estimated costs of fishing. Commercial viability is subjectively determined by the Scallop Fishery Advisory Committee (ScFAC), which is largely comprised of industry members.
3. Listing candidate open areas – The spatial management regime requires areas to be assessed to characterize population structure and extent of candidate scallop beds before being considered for commercial harvesting. Initially, statewide exploratory surveys are conducted using industry scallop vessels that can retain limited tonnages of located scallops (50 t survey area cap for six areas; 300 t State-wide, see Figure 7). If targeted surveys with an observer on board then reveal that the size structure indicates a discard rate no larger than 20% prior to an opening, then the surveyed area can be considered as a candidate area for a fishery opening. The effect of the spatial management regime is essentially a rotational harvest system.
4. Discard threshold – If a vessel has more than two dredge tows in which $\geq 20\%$ of scallops are less than minimum size, then the vessel should move at least 250 m. If such discard rates persist, then the coordinates should be reported for potential mitigation measures.
5. Meat recovery guideline – As a general guideline, meat recovery (adductor muscle and roe) should be >11.8 g corresponding to <85 scallops/kg.

Table 2: Summary of the Tasmanian scallop fishery

Area	The area of the fishery extends from the high-water mark to 20 nautical miles into Bass Strait and from the high water mark out to 200 nautical miles off the rest of the State of Tasmania.
Fishery status	The target species within Tasmanian waters is currently considered to be overfished.
Target Species	Commercial scallops (<i>Pecten fumatus</i>).
By-product Species	Doughboy (<i>Mimachlamys asperima</i>). Queen scallop (<i>Equichlamys bifrons</i>).
Gear Commercial	Benthic scallop dredge.
Recreational/Indigenous	Dive only – D'Entrecasteaux Channel closed to fishing.
Season	Peak catch and effort occur between winter and spring/early summer (Fishery closes 31 December).
Primary landing ports	Stanley, Triabunna, St Helens.
Fishing licences	62
Active vessels	2 in 2021 season. Most operators also have access to one or both of the Victorian and Bass Strait Central Zone (BSCZ) scallop fisheries.
Value of commercial harvest	Up to \$6 million, with the 2004 beach value of scallops being \$5.5 million.
Management arrangements	Output controlled through: • TAC based on aggregation of 10,366 scallop bag units, • the Minister can set and alter the quota unit value by public notice. Input controlled through: • limited entry (fishers must also hold a scallop entitlement) and a minimum unit holding to operate, • minimum size limits to allow for two spawning's, • spatial management regime, where most of the fishery area is closed and only certain defined areas opened if criteria met, • seasonal closure – fishing only allowed when scallops have reached optimum condition and to maximise recruitment, • limits on number, dimensions and structure of dredges, • 1 April fishery opens as state-wide survey, survey period to end when an open season is declared or on 31 December.
Export	Mainly a domestic market, although export grew substantially in 2005, particularly to France.
Bycatch	Bycatch is relatively low, consisting mostly of molluscs such as dog cockles (<i>Glycymeris</i> sp.) and the native oyster (<i>Ostrea angasi</i>). Diogenid hermit crabs (<i>Paguristes tuberculatus</i>) and the introduced screw shell (<i>Maoricolpus roseus</i>) are also taken as bycatch.
Interaction with Threatened Species	Considered low, but potentially greater interactions with syngnathids. Possible minor interactions with seals, sharks, cetaceans and seabirds.




Figure 7. State-wide survey areas.

Recreational fishery

Recreational fishing for scallops has a long history in Tasmanian waters and, like the commercial fishery, has been characterised by open seasons followed by extended closures for stock rebuilding. The current recreational dive-only fishery for scallops (commercial, queen, and doughboy scallops are targeted) commenced in 2006 and catches are input controlled with open seasons, closed areas, size limits, bag limits, possession limits, and a specific scallop diver recreational license. Although the entire Tasmanian coast was initially open to recreational effort, TAFI (now known as IMAS) recreational fishing surveys showed that fishing was primarily focused in the D'Entrecasteaux Channel, and consequently IMAS conducted seven scallop dive surveys of the channel between 2006 and 2011. This research established that the total number of scallops in the channel declined by 87% between 2006 and 2011 (commercial scallops declined by 93%). These surveys and subsequent D'Entrecasteaux Channel surveys in 2012 (dive), 2016 (dive), 2017 (video and dive) and 2020 (video) found that no significant recruitment pulses had occurred since 2007. On this basis, the D'Entrecasteaux Channel component of the recreational fishery has been closed to fishing since 2012 and is likely to remain closed for the foreseeable future due to low abundance of juveniles.

The recreational scallop dive fishery elsewhere around the state has supported open seasons each year from 2012 and approximately 13,000 and 130,000 scallops landed in the 2013 and 2018 recreational seasons respectively, primarily from the central east coast of Tasmania (Lyle *et al.*, 2014; Lyle *et al.*, 2019).

Basic commercial scallop biology

Commercial scallops (*Pecten fumatus*) are filter feeders that sift plankton and detritus from the water column. They are simultaneous hermaphrodites and highly fecund, with up to 1 million eggs produced by an individual scallop. Reproduction is by broadcast spawning, where individuals release sperm first followed by eggs, (Minchin, 2003). In Tasmania, *P. fumatus* has a protracted spawning season involving several partial spawning events, as spawning lasts 5–6 months during spring and summer (Mendo *et al.*, 2014a). After external fertilization, larvae remain in the water column for 30 d before settling on fine to coarse sand (generally without organic sediment) forming aggregations commonly referred to as “beds” (Mendo *et al.*, 2014a). The longevity of scallop beds depends on cohort structure. If beds consist of a single cohort, the whole bed disappears when the animals reach the end of their lifespan (Hortle and Cropp, 1987). The size of scallop beds can vary from 5 to 30 square nautical miles (17–100 km²) (Haddon *et al.*, 2006). The settled scallops grow quickly and reach 70 to 75 mm shell length (SL) in around 18 months. Age and size at maturity are 2 years and 70-80 mm SL, and maximum age and size are 7 years and 120mm; however, these parameters are known to vary by region.

2. Fishery peer review

A peer review of the scallop assessment process was conducted by Dr Gordon H. Kruse in 2015 (Kruse, 2015). Dr Kruse was employed from 1985-2001 at the Alaska Department of Fish and Game, first as the Statewide Shellfish Biometrician and then as Chief Marine Fisheries Scientist; and from 2001 to present has been the President's Professor of Fisheries at the University of Alaska Fairbanks. With respect to scallops, he led the development of the State of Alaska's scallop fishery management plan and co-authored the first federal scallop fishery management plan for Alaska. He served on the Scallop Plan Team of the North Pacific Fishery Management Council for 8 years and is in his 15th year as a member on the Scientific and Statistical Committee of the North Pacific Fishery Management Council, where he conducts reviews of the annual Stock Assessment and Fishery Evaluation (SAFE) reports for scallops and other species. Recently, he conducted a scientific review of the weathervane scallop fishery in Alaska for Monterey Bay Aquarium's Seafood Watch Program. In addition to the state and federal management plans for scallops, he has published 11 peer-reviewed papers and agency reports on the scallop fishery in Alaska.

The review was informed by interviews with industry, researchers and fishery managers and through documents including scientific papers, survey reports, and minutes of advisory committee meetings relating to application of research data for decision-making. The review was first prepared as a draft, and then provided to Government and Institute for Marine and Antarctic Studies (IMAS) to enable any correction of factual errors or oversights. The aims of the review were:

1. Evaluate the adequacy of life history information delivered by the current assessment program to fishery assessment and management decision-making processes.
2. Evaluate the adequacy of monitoring data used for assessing current status of the stock and used for informing management decisions and the annual harvest strategy (i.e., open areas and Total Allowable Catch, TAC, setting).
3. Evaluate the adequacy and application of methods used to translate life history and monitoring data into scientific guidance on the state of the stock and harvest scenarios.

Reviewer's Recommendations

The reviewer's recommendations were in three categories, as detailed below. Text in *italics* outlines how this has been addressed:

1. Improved understanding of *Pecten fumatus* life history through research into:
 - Stock structure using genetic and larval drift methods

Addressed in FRDC 2008/022: Establishing fine-scale industry based spatial management and harvest strategies for the commercial scallop fishery in Southeast Australia (see page 14)

- Reproductive biology

Addressed in FRDC 2008/022 and FRDC 2012/027: Determining when and where to fish: Linking scallop spawning, settlement, size and condition to collaborative spatial harvest and industry in-season management strategies (see page 11).

- Age, growth and maturity

There is currently no simple reliable and repeatable technique for accurately ageing scallops. As such, age is currently estimated from a previously defined shell length to age relationship.

Growth has been examined in FRDC 2003/017: Juvenile scallop trashing rates and bed dynamics: testing the management rules for scallops in Bass Strait, using modal analysis of length frequency data (see page 15) and was further addressed using this technique in FRDC 2012/027.

The current monitoring program is generally adequate to inform the current paddock fishing strategy, although future improvement could include allocation of size limits specific to growth rates by area.

A lower minimum size limit (85 mm SL) is applied to scallop beds in the north-west bed given the historically slow growth rates in this region and the fact that most scallops of this size are still 3+ years old, and hence have had two major spawning's (Martin et al., 1988).

2. Improved scientific guidance to management of the fishery through provision of regular stock assessment reports including:
 - Historic landings: *See pages 4-10.*
 - Survey results including size frequency distributions: *Survey undertaken in 2020 identifying size frequency distributions and legal sized density across fishery reported by Ewing and Semmens (2020).*
 - Catch per unit effort (CPUE) reporting: *Fishery closed since 2015.*
 - Economic performance of the fishery: *Fishery closed since 2015.*

3. 2021 Pre-season survey

Introduction and survey methods

The 2020 Tasmanian Scallop Fishery (TSF) preseason assessment yielded scallop density and size structure data over a wide spatial scale. However, it also inadvertently resulted in significant dredge effort expended on scallop beds with high discard rates to utilize the allowable retained catch component of the survey. Consequently, the rules for participation in the 2021 TSF preseason dredge survey were reformulated to ensure that significant survey dredge effort would not be expended on beds with high discard rates. The three categories of the preseason dredge survey, opportunistic, exploratory, and targeted surveys, were retained in the new system.

Dredge surveys

The 2021 (TSF) pre-season assessment was comprised of only targeted commercial dredge surveys and IMAS video surveys (see Ewing et al., 2021 and Semmens et al., 2023 for details). Regions identified in the 2020 survey as known scallop beds (White Rock, The Gardens, and East Flinders) were demarcated and were not available for opportunistic or exploratory dredge surveys. IMAS developed a random survey design within these regions for the targeted component of the dredge survey to better reflect the variation in density of scallops across the defined area of interest. Fishers participating in targeted dredge surveys were chartered by DPIPWE to sample in accordance with the IMAS random survey design and did not retain scallops. The sites towed were randomly assigned within the boundaries of polygons defined by DPIPWE managers as the regions of commercial interest (Table 3, Figure 8). An IMAS or department observer was onboard for targeted surveys and catch weights (kg) of legal-sized scallops (> 90 mm) were recorded for each tow. The observer recorded the size structure of each tow by measuring the shell length (width at the widest point of the shell) of up to 100 scallops randomly sampled from the ungraded catch. The density of legal-sized scallops was reported as the ratio of the landed weight of legal sized scallops over the product of dredge width and tow length (calculated from start and finish coordinates).

Video Surveys

The video component of the survey was conducted in the White Rock region of interest in June 2021 (Table 3). Forty-eight sites within this region were randomly assigned for the 2020 White Rock Video Survey using the Create Random Points tool in the software package ArcGIS 10.4. Assignment of random points was constrained within a 200 km² polygon (perimeter 72 km) provided by DPIPWE that defined the area of interest to management, and with a minimum allowable separation of 500m. The 2021 survey deployed the IMAS towed video camera unit at the sites from the 2020 video survey and towed from the site position in a straight line transect for at least 500m, approximately 1m above the seafloor, and at around 1.5 knots. Each transect was recorded as a track on the vessel GPS unit.

Video footage from each site (in a randomised order) was viewed and analysed using video analysis software Transect Measure (SeaGIS) which allows frame-by-frame analysis of the video footage. Within any frame a pixel-to-millimetre calibration was made using a known distance (e.g., the separation of scaling lasers) allowing objects within that frame to be measured in mm. The software also incorporated annotation of the position and time sequence of objects and measurements within any frame. By matching the frame time sequence with vessel GPS track time sequence data, the geographic position of objects in a frame was determined. These features of the software were used to ascertain the number of commercial scallops, their size, and their geographic position within the analysed segment of each video transect.

Table 3: Survey tows conducted in the 2021 pre-season survey by region and survey method.

Area	Region	Dredge			Video	
		Vessel	Targeted	Exploratory		Opportunistic
3	North Babel	Dell Richey II	30	-	-	-
	South Babel	Dell Richey II	30	-	-	-
4	Pot Boil	Dell Richey II	30	-	-	-
	The Gardens	Dell Richey II	28	-	-	-
6A	White Rock	Odete C	29	-	-	48
		Total	147			48




Figure 8. Map showing dredge survey tows (black dots).

Survey results by region

Area 6A White Rock

The White Rock targeted dredge survey consisted of 29 survey dredge tows deployed from FV Odete C in May 2021 (Table 3; Figure 8). The targeted dredge sites were a random subset of the 48 sites randomly allocated in the White Rock region for the 2020 White Rock Video survey. The 2021 White Rock video survey consisted of 48 tows of the IMAS towed video unit. Video tows were each of around 500m in length and were also deployed on the sites from the 2020 White Rock Video survey.

White Rock targeted dredge survey tows yielded an average density of legal-sized scallops of 15.6 kg/1000 m² (median 14.2 kg/1000 m²), with a discard rate of 5.67% (Table 4; Figure 9A). Results from the video survey yielded an average density of legal-sized scallops of 30.4 kg/1000 m² (median 27.2 kg/1000 m²), with a discard rate of 30.5% (Table 4; Figure 9B). Higher representation of smaller scallops in the video survey is expected and reflects the size selectivity of commercial scallop dredges.

The low to moderate densities of legal-sized scallops in the 2021 survey were similar to those encountered in the 2020 pre-season survey. The lower proportions of sub-legal scallops and higher average sizes relative to the 2020 survey reflect growth into the fishery of the standing crop of scallops but may also indicate reduced stock recruitment. The full video survey results may provide more clarity on size structure when available.

A low dredge survey discard rate, a decline in the video survey discard rate and high average size indicate that the White Rock region has significantly less undersize scallops in 2020. However, given the lack of recent recruitment, low to medium densities of legal-sized scallops and the past history of openings and closures in this area, the merits of opening White Rock in 2021 need to be given careful consideration, noting the depleted status of the Tasmanian scallop fishery.

Table 4. Legal-sized scallop density, average size, and discard rate for the 6A White Rock sub-region from recent dredge and video pre-season surveys.

Year	Method	Sub-Area 6A			
		Density (kg/1000 m ²)		Mean scallop size (mm ± SE)	Discard rate (% <90 mm)
		Mean	Median		
2021	Dredge	15.6	14.2	102.2 ± 0.2	5.7
	Video	30.4	27.2	96.0 ± 0.4	30.5
2020	Dredge	20.9	16.6	97.9 ± 0.2	7.5
	Video	21.9	8.5	85.6 ± 0.4	58.9
2017	Dredge	4.7	3.9	100.5 ± 0.1	2.2
2016	Dredge	4.4	3.5	96.8 ± 0.2	22.5
2015	Dredge	19.7	15.8	99.1 ± 7.7	3.9
2014	Dredge	8.6	7.2	98.2 ± 8.2	7.3
2013	Dredge	6.2	-	93.9 ± 11.1	18.2
2011	Dredge	Dredge	>30	-	-




Figure 9. Legal-sized scallop density per tow for the White Rock dredge targeted survey (A) and from the video survey (B). Density is displayed as kg (of scallops >90mm) per 1000 m².




Figure 10. Standardised length frequency distribution for the 29 dredge tows in the White Rock targeted dredge survey (A) and video survey results (B). The dredge discard rate was 5.67% from 2413 scallops measured at 90 mm minimum length with a mean size $\pm$ se: 102.2 ± 0.2 mm. The video discard rate was 30.5% from 1220 scallops measured at 90 mm minimum length with a mean size $\pm$ se: 96 ± 0.4 mm. The vertical lines show the 90 mm minimum length.

The Gardens

The Gardens targeted survey consisted of 28 survey dredge tows deployed from FV Dell Richey II in June 2021 (Table 3; Figure 8). The Gardens survey tows yielded an average density of legal-sized scallops of 12.3 kg/1000 m² (median 4.8 kg/1000 m²), with a discard rate of 61.3% (Table 5; Figures 11 and 12). Average densities of legal-size scallops were similar to those encountered in the non-random 2020 survey. Both surveys yielded very low median densities indicating that average densities were strongly influenced by a small number of sites with relatively high scallop densities.

Very high discard rates and patchy distribution indicate that the Gardens is currently not suitable for opening (Table 5; Figures 11 and 12).

Table 5. Scallop density, average size, and discard rate for The Gardens region in Area 4 in 2020 and 2021 pre-season surveys.

Season	Density (kg/1000 m ²)		Average Scallop size (mm ± SE)	Discard % <90mm
	Average	Median		
2021	12.3	4.8	90.5 ± 0.3	61.3
2020	16.9	4.2	100.8 ± 0.4	33




Figure 11. Legal-sized scallop density per shot for The Gardens targeted survey. Density is displayed as kg (of scallops >90mm) per 1000 m².




Figure 12. Standardised length frequency distribution for the 28 tows in the targeted survey conducted in The Gardens (Area 4). Discard rate was 61.3% from 1563 scallops measured at 90 mm minimum length. Mean size $\pm$ se: 90.5 $\pm$ 0.32 mm. The vertical line shows the 90 mm minimum length.

Pot boil

The targeted survey conducted in the Pot Boil region to the east of Flinders Island in Area 4 consisted of 30 survey dredge tows deployed from FV Dell Richey II in June 2021 (Table 3; Figure 8). The Pot Boil survey tows yielded an average density of legal-sized scallops of 20.9 kg/1000 m² (median 19.1 kg/1000 m²), with a discard rate of 43.6% (Table 6; Figures 13 and 14).

Very high discard rates and moderate densities encountered at the Pot Boil in both the 2020 non-random and 2021 random surveys, indicate that this region is currently not suitable for opening (Table 6, Figures 13 and 14).

Table 6. Scallop density, average size, and discard rate for the Pot Boil region in Area 4 in 2020 and 2021 pre-season surveys.

Season	Density (kg/1000 m ²)		Average Scallop size (mm $\pm$ SE)	Discard % <90mm
	Average	Median		
2021	20.9	19.1	92.2 $\pm$ 0.18	43.6
2020	19.7	14.9	93.1 $\pm$ 0.24	35.9




Figure 13. Legal-sized scallop density per shot for the Pot Boil region in Area 4 targeted survey. Density is displayed as kg (of scallops >90mm) per 1000 m².




Figure 14. Standardised length frequency distribution for the 30 tows in the targeted survey conducted in Pot Boil region (Area 4). Discard rate was 43.6% from 2069 scallops measured at 90 mm minimum length. Mean size $\pm$ se: 92.2 $\pm$ 0.18 mm. The vertical line shows the 90 mm minimum length.

South Babel

The targeted survey conducted in the South Babel region to the east of Flinders Island in Area 4 consisted of 30 survey dredge tows deployed from FV Dell Richey II in June 2021 (Table 3; Figure 8). The South Babel survey tows yielded an average density of legal-sized scallops of 38.3 kg/1000 m² (median 36.3 kg/1000 m²), with a discard rate of 26.7% (Table 7; Figures 15-17). While the densities of legal-sized scallops were distributed evenly throughout the South Babel region (Figure 15), discard rates were low in the north (Figures 16 and 17B), and high in the south of the region (Figures 16 and 17C). Consequently, average densities have also been provided for the upper and lower portions of South Babel region (Table 7).

Table 7. Scallop density, average size, and discard rate for the South Babel region in Area 4 in 2020 and 2021 pre-season surveys. Values have also been provided for the subsets of the region above and below Latitude 40° south in the 2021 pre-season survey.

Season	Sub-Region	N tows	Density (kg/1000 m ²)		Average Scallop size (mm $\pm$ SE)	Discard % <90mm
			Average	Median		
2021	Total	30	38.3	36.3	96.6 $\pm$ 0.22	26.7
2021	Nth of 40° S	20	35.6	32.6	98.6 $\pm$ 1.22	13.4
2021	Sth of 40° S	10	40.1	33.0	92.0 $\pm$ 0.38	39.7
2020	-	32	27.4	25.0	94.4 $\pm$ 0.21	29.3




Figure 15. Legal-sized scallop density per shot for the South Babel region in the Area 4 targeted survey. Density is displayed as kg (of scallops >90mm) per 1000 m².




Figure 16. Discard percentage per shot for the South Babel region in the Area 4. Percentage text values indicate standardised discard rates for the portion of the South Babel region above and below Latitude -40° south.




Figure 17. Standardised length frequency distributions for the targeted survey conducted in South Babel region (Area 4). Plot A shows the distribution for the whole region (30 tows, 26.7% discard); plot B shows the distribution for sites that lie in the portion of the region north of Latitude 40° south (20 tows 13.4% discard); and plot C shows the distribution for sites that lie in the portion of the region south of the Latitude 40° south (10 tows 39.7% discard). Mean size $\pm$ se for plot A was 96.62 ± 0.22 mm, for plot B was 98.63 ± 1.22 mm, and for plot C was 92.04 ± 0.38 mm. The vertical line shows the 90 mm minimum length.

North Babel

The targeted survey conducted in the North Babel region to the east of Flinders Island in Area 3 consisted of 30 survey dredge tows deployed from FV Dell Richey II in July 2021 (Table 3; Figure 8). The North Babel survey tows yielded an average density of legal-sized scallops of 24.1 kg/1000 m² (median 19.8 kg/1000 m²), with a discard rate of 5.4% (Table 8; Figures 18 and 19).

Low discard rates and moderate densities were encountered at North Babel in both the 2020 non-random and 2021 random surveys (Table 8, Figures 18 and 19). This area meets the harvest strategy rules for opening. However, this is a relatively small area, with a relatively small biomass (Tables 8 and 9), and as such may not be suitable to opening on its own.

Table 8. Scallop density, average size, and discard rate for the North Babel region in Area 3 in 2020 and 2021 pre-season surveys.

Season	Density (kg/1000 m ²)		Average Scallop size (mm ± SE)	Discard % <90mm
	Average	Median		
2021	24.1	19.8	103.8 ± 0.29	5.4
2020	24.0	13.2	103.8 ± 0.24	3.73




Figure 18. Legal-sized scallop density per shot for the North Babel region in the Area 3 targeted survey. Density is displayed as kg (of scallops >90mm) per 1000 m².




Figure 19. Standardised length frequency distribution for the 30 tows in the targeted survey conducted in North Babel (Area 3). Discard rate was 5.4% from 1409 scallops measured at 90 mm minimum length. Mean size $\pm$ se: 103.8 ± 0.29 mm. The vertical line shows the 90 mm minimum length.

Biomass estimates

An estimated biomass of scallops within each of the polygons defined by DPIPWE managers as the regions of commercial interest was calculated using the standard method of scaling-up the mean density across the survey area for both the dredge and video surveys (Table 9). Biomass estimates have also been provided for the upper and lower portions of South Babel region because, despite similar densities, discard rates were low in the north of the region and high in the south of the region (Table 9). Note that adjusting the boundaries of the polygon containing the randomly allocated shots after conducting the survey violates the assumptions made in the analysis and as such the biomass values from the two new regions (North and South sections of the larger South Babel region) should be assessed with relative caution.

An adjustment for dredge efficiency of 33% (Harrington et al. 2008b) has been applied to the dredge biomass estimates and is also presented in Table 9. It should, however, be noted that the comparison between the dredge and video biomass estimates in the White Rock region in the 2021, and 2022 surveys raises questions around the validity of the adjustment applied for dredge efficiency. The dredge estimates of the mean and lower and upper 95% confidence intervals for biomass were 61.3, 63.4 and 60.3% of those for video in 2021, respectively. Similarly, the dredge estimates of the mean and lower and upper 95% confidence intervals for biomass were 58.1, 54.3 and 60.2% of those for video in 2022. This suggests that dredge efficiency, at least in the White Rock region, is significantly higher than 33% and closer to 60%. Consequently, dredge biomass values provided here should only be used as a relative value for comparison between the regions surveyed.

Table 9. Density and estimated biomass of legal-sized commercial scallops (*Pecten fumatus*) within each of the polygons defined by DPIPWE managers as the regions of commercial interest, based on dredge and video survey methods. Biomass estimates are also provided for the portions of the South Babel region that lie above and below the latitude 40° south. Dredge-derived biomass estimates adjusted for dredge efficiency should be interpreted with caution as recent White Rock video surveys raise questions about their validity. Biomass confidence intervals are 95%.

Region	Sub-region	Method	n	Sub-legal %	Mean kg/m ²	SD kg/m ²	Area (km ²)	Biomass (t)			Adjusted Biomass (t)		
								lower CI	Mean	higher CI	lower CI	Mean	higher CI
White Rock	-	Dredge	29	5.7	0.0156	0.0144	200	2025	3122	4220	6072	9366	12660
	-	Video-2021	38	30.5	0.0304	0.0286	200	3191	5093	6995	-	-	-
	-	Video-2020	48	58.9	0.0219	0.028	200	2077	3161	4246	-	-	-
Gardens	-	Dredge	28	61.3	0.0123	0.0184	10.4	54	128	202	162	384	606
Pot Boil	-	Dredge	30	43.6	0.0210	0.0136	16.4	260	343	426	780	1029	1278
South Babel	-	Dredge	30	26.7	0.0371	0.0246	67.6	1888	2509	3131	5664	7527	9393
South Babel	Nth 40°S	Dredge	20	13.4	0.0356	0.0200	40.9	1072	1456	1839	3216	4368	5517
South Babel	Sth 40°S	Dredge	10	39.7	0.0401	0.0330	26.7	422	1073	1705	1266	3219	5115
North Babel	-	Dredge	30	5.4	0.0241	0.0212	13.9	224	334	444	672	1002	1332

Conditions for opening the 2021 commercial scallop season

The Pot boil and the Gardens showed high discard rates of 43.6 and 61.3 % respectively, and patchy distribution and are not recommended for opening in 2021 (see Table 10). Low discard rates and moderate densities were encountered at North Babel in both the 2020 non-random and 2021 random surveys. This area meets the harvest strategy rules for opening. However, this is a relatively small area, with a relatively small biomass, and as such may not be suitable to opening on its own. A small area south of 40°S in the South Babel region meets the requirements for opening in 2021. White Rock has shown a lack of recent recruitment and low to medium densities of legal-sized scallops. Given the past history of openings and closures in this area, the merits of opening White Rock in 2021 need to be given careful consideration, noting the depleted status (status in 2020) of the Tasmanian scallop fishery.

Table 10. Summary of the surveys conducted during the 2021 pre-season survey.

Area	Region	Dredge/Video Tows	Ave Density kg/1000m	Average Size (mm ± SE)	Discard % <90 mm
3	North Babel	30	24.1	103.8 ± 0.3	5.4
	South Babel total	30	38.3	96.6 ± 0.2	26.7
	South Babel Nth of 40° S	20	35.6	98.6 ± 1.2	13.4
	South Babel Sth of 40°S	10	40.1	92.0 ± 0.4	39.7
4	Pot Boil	30	20.9	92.2 ± 0.18	43.6
	The Gardens	28		90.5 ± 0.3	61.3
6A	White Rock Dredge	29	15.6	102.2 ± 0.2	5.7
	White Rock Video	48	30.4	96.0 ± 0.4	30.5

4. Economic data

The TSF was partly opened in 2021, the first season to open since 2015, with the TAC set at 1494.9 tonnes or 145 kg/unit. The open areas consisted of reporting blocks 3H30 and 3H3P in the vicinity of Babel Island east of flinders island, it excludes waters less than 20 m and all commonwealth waters (Figure 20). Only 44.5 tonnes were landed due to poor quality and reported decrease in recoveries by processors. The remainder of the state continued to be closed.

In 2021, there were 62 scallop licenses held by 62 quota holders, with a total of 10336 quota units. In 2021, there were only three processors that received scallops.




Figure 20. Map of 2021 scallop commercial harvest area, (i) 39° 52.5'S, 148° 15'E; (ii) 39° 52.5'S, 148° 30'E; (iii) 40° 00'S, 148° 30' E; (iv) 40° 00'S, 148° 15'E.

5. Warm water event

During July 2021 processors and the fishery reported a drop in recoveries, affecting both Commonwealth and state beds from Lakes Entrance to Flinders Island. Fishers reported a rise in sea surface temperature of up to 1.5 °C. Subsequently, when Tasmanian Scallop Area 3 opened to fishing in September 2021, there had been a major die-off post survey, most likely due to the warm-water incursion in the Babel Island area in July (e.g., see Figure 21).




Figure 21. SST data showing a warm water incursion around Babel Islan in July 2021.

6. Stock status

STOCK STATUS	RECOVERING
The Tasmanian Scallop Fishery (TSF) is managed with a harvest strategy where surveys are undertaken to estimate abundance and decision rules are used to open areas to fishing with total allowable catches (TACs) based on the estimated abundance. Biomass in the Tasmanian Scallop Fishery (TSF) is historically overfished (Caton and McLoughlin, 2004), with recruitment and production levels now affected. In 2013, 2014 and 2015, surveys generally found low scallop densities and limited evidence of successful recent recruitment but did identify two beds (one on the north-west coast and the other on the east coast) containing commercial quantities (Ewing <i>et al.</i>, 2016). Surveys in 2016 and again in 2017 generally only found very low levels of scallop abundance and limited evidence of successful recruitment, with no area considered to contain commercially viable quantities in either year (Ewing <i>et al.</i>, 2017; Semmens <i>et al.</i>, 2018). This includes the east and north-west coast beds fished in 2013-2015, which appeared to have been fished down to a commercially unviable density, with no subsequent recruitment evident. Given the results of the 2016 and 2017 surveys, there was a low expectation that conducting pre-season surveys in 2018 and 2019 would yield the presence of commercially viable scallop beds, and as such, pre-season surveys were not conducted. In 2020 a pre-season survey was conducted, and the survey results were dominated by low to moderate densities of legal sized scallops, several areas demonstrated signs of recruitment that suggest the possibility of supporting a tightly managed commercial fishery in the near- to medium-term. The 2021 pre-season survey consisted of 147 targeted dredge tows and 48 video surveys. Survey results showed low to moderate densities of legal sized scallops, with patchy high densities and variable discard rates. Fishing mortality is managed with the aim of restricting catches to beds of mature scallops near the end of their lifespan. The combination of the harvest strategy and depleted biomass has led to a history of closures due to low abundance. In recent times, the fishery was closed between 2000-2002 and again between 2009 and 2010. Areas with commercial density of scallops towards the end of their lifespan were opened to fishing each year between 2013 and 2015. The fishery was closed between 2016 and 2020. The TSF was partly opened in 2021, the first season to open since 2015, with the TAC set at 1494.9 tonnes or 145 kg/unit. However, only 44.5 t were landed from the open area due to a major die-off event post survey. The remainder of the state continued to be closed. On the basis that biomass has increased during the 2016-20 closure due to wide-scale recruitment events, and current restrictions are effectively limiting fishing mortality, the TSF is classified as recovering.	
STOCK	Tasmanian Scallop Fishery
INDICATORS	Size structure, density and catch

7. Bycatch and protected species interaction

Bycatch and protected species interaction from the recreational fishery are effectively nil as scallops are hand collected. Vessels usually anchor at dive locations but anchors are deployed on unconsolidated sediment so there is no concern with habitat interaction.

Bycatch impacts of the commercial fishery are also low because volume is low, and almost all bycatch is returned alive and healthy. Common bycatch species are molluscs such as dog cockles (*Glycymeris* sp.) and the native oyster (*Ostrea angas*). Diogenid hermit crabs (*Paguristes tuberculatus*) and the introduced screw shell (*Maoricolpus roseus*) are also taken as bycatch (see a more detailed list from scientific analysis of scallop dredge catches at White Rock in 2006, Table 11).

Interactions with threatened, endangered or protected species (TEPS) are rare. There are low levels of interactions with syngnathids (seahorses and related species) observed, but these are usually released unharmed and interactions with syngnathid populations are limited, given that most of the fishing effort takes place in habitats that are not generally favoured by syngnathids (i.e., sandy bottoms with low levels of habitat structure). Gear is towed at low speed so interactions with mobile protected species like seals, sharks, cetaceans and seabirds are not observed.

When the fishery is opened, scallop fishers need to record any interactions with TEPS, and record retained by-product, no reports were received in 2021. Fishers are only authorised to retain scallop species, so by-product only consists of queen (*Equichlamys bifrons*) and doughboy (*Mimachlamys asperima*) scallops, of which none were retained in 2021. Fishers do not record bycatch caught in the dredge. Scallop fishers participating in statewide surveys may make notes and observations about bycatch in log sheets, but this is not compulsory. For targeted surveys, where there are observers on board, the observers (IMAS or DPIPWE staff) may make some more detailed observations, including any interactions with TEPS, but the bycatch from each dredge tow is not quantified. More detailed collection of bycatch information during surveys should be considered, given that these are the only means of collecting this data for the fishery.

Habitat interaction from the commercial fishery has been of some concern in the past when gear interacted with sponge habitat. This problem is now overcome with the system of smaller spatial boundaries for beds opened in the TSF. This process takes account of sensitive habitats to ensure no gear interaction.

Table 11. List of scallop bed species retained in commercial gear during the two scallop dredge surveys conducted at White Rock in 2006.

Category	Common Name	Scientific Name
Commercial Scallop	Commercial Scallop	<i>Pecten fumatus</i>
Bivalves	Doughboy Scallop	<i>Mimachlamys asperima</i>
	Queen Scallop	<i>Equichlamys bifrons</i>
	Mud Oyster	<i>Ostrea angasi</i>
	Mussel	<i>Mytilus edulis</i>
	Dog Cockle	<i>Glycymeris striatularis</i>
	Razor clam	<i>Atrina tasmanica</i>
Other Molluscs	New Zealand Screw Shell	<i>Maoricolpus roseus</i>
	New Holland Spindle Shell	<i>Fusinus novaehollandiae</i>
	Tulip Shell	<i>Pleuroploca australasia</i>
	Triton Shell	<i>Charonia lampas</i>
	Cowrie	<i>Cypraea hesitata</i>
	Pale Octopus	<i>Octopus pallidus</i>
Crustaceans	Southern Keeled Octopus	<i>Octopus berrima</i>
	Hermit Crab	<i>Strigopagurus strigimanus</i>
	Unidentified Hermit Crab	Unidentified sp.
	Spider Crab	<i>Leptomithrax gaimardii</i>
	Hairy Shore Crab	<i>Pilumnus tomentosus</i>
	NZSS Hermit Crabs	Unidentified spp.
Seastars	11-Arm Seastar	<i>Coscinasterias muricata</i>
	Astropectinid	<i>Bollonaster pectinatus</i>
	Oreasterid	<i>Nectria ocellata</i>
Urchins	Common Urchin	<i>Heliocidaris erythrogramma</i>
	Pencil Urchin	Unidentified sp.
Fish	Crested Flounder	<i>Lophonectes gallus</i>
	Lachet	<i>Lepidotrigla vanessa</i>
	Shaw's Cowfish	<i>Aracana aurita</i>
Rays	Tasmanian Numbfish	<i>Narcine tasmaniensis</i>
	Banded Stingaree	<i>Urolophus cruciatus</i>
Other Species	Pumpkin Sponge	Unidentified sp.

8. Conclusions and recommendations

The 2021 preseason survey suggested that scallops remained in low to moderate densities around most of the Tasmanian coast. High discard rates were found at Pot Boil and the Gardens along with patchy scallop distributions. Small patchy areas of higher density were identified around Babel Island (both North and South sites).

FRDC 2008-022 (Mendo *et al.* 2014b; Semmens *et al.* 2015) demonstrated that individual scallops are more likely to spawn when at higher densities and at closer proximity to other scallops, with the potential benefits of increasing encounter rates between gametes (eggs and sperm), and the likelihood of spawning success. Maintaining regions of high scallop density is, therefore, considered an important strategy to enhance recruitment success, partly due to the improved fertilization rates at greater densities, but also due to the indirect effect on synchronization of spawning, i.e., more scallops spawn at the same time when there are more of them, and they are closer to each other (higher density beds).

There were relatively small areas of higher density scallops in some regions of the fishery, although they are generally spatially separated (i.e., there were no large beds with high densities). Of the beds sampled in the 2021 preseason survey, patches of higher densities were found at both North Babel and South Babel (average of 24.1 and 38.3, and median of 19.8 and 36.3 kg/1000 m², respectively). There were, however, large differences in the discard rate, with North Babel having a discard rate of 5.4% and South Babel 26.7%.

The depleted nature of the Tasmanian commercial scallop fishery and the recent closure (2016-2020) following the likely premature opening of the fishery in 2015, suggested that the precautionary principle needed to be applied in the absence of further information. As such only a limited open of the fishery was undertaken in 2021 in Area 3 (1494.9 tonnes).

However, during July 2021 processors and fishers reported a drop in scallop meat recovery (meats/kg), affecting both Commonwealth and state beds from Lakes Entrance to Flinders Island. Fishers reported a rise in sea surface temperature of up to 1.5 °C. Subsequently, when Tasmanian Scallop Area 3 opened to fishing in September 2021, there had been a major die-off post survey, most likely due to a warm-water incursion in the Babel Island area in July. Sudden bed die-off events in the eastern portion of the fishery may become more common with warming waters from the east Australian current pushing further southward and predicted heatwave events. This will need to be factored into the planning and management of the fishery.

The survey scheme used in and prior to 2020 was conceived before the extended fishery closure between 2016-2020 and was a practical tool for achieving surveys at that time. However, its shortcomings are amplified in a depleted/recovering fishery. Although the survey tows conducted in the survey scheme in 2020 were unlikely to impact density and hence potential recruitment, they were accompanied by targeted fishing associated with the survey quota allocation of 50 t/survey area (Figure 7). For example, the 2020 survey entailed 635 dredge tows and over 1300 targeted fishing shots. This targeted fishing occurs in the densest areas found during the survey, reducing the density of the beds, which may in turn impact potential recruitment in a depleted fishery, prior to a decision being made as to whether the biomass can support a fishery and what areas should be closed due to undersized scallops. This effect was pronounced in Area 4 during the 2020 preseason survey where discard rates were particularly high (up to 44%) in higher density beds likely to be targeted to collect the survey quota. It should be noted that the survey scheme was adjusted in 2021 to limit the impact of targeted fishing and time will tell if the new scheme does in fact achieve this aim.

Along with targeted fishing, the survey scheme used in and prior to 2020, did not use randomly designated dredge tows. The data from the 2020 survey showed that this was leading to the fishers sometimes clumping survey shots around legal sized scallops, rather than fully surveying the area, or alternatively not fully investigating an area when legal sized scallops were not initially found. Both these unconscious behaviours limited the effectiveness of the surveying. It should be noted that the survey scheme was adjusted in 2021 to encompass a random survey design. Unlike the

clumped/limited sampling in the 2020 survey, the random sampling method better reflects the variation in density of scallops across the defined area of interest.

Video surveys offer an advantage over dredge surveys due to: (i) lower costs for data collection; (ii) enhanced precision of size structure; (iii) information on recruitment in recent years; (iv) minimal impact on the seafloor and benthic biota and (v) no mortality of sub-legal scallops. Their continued use is recommended for providing data for management of the TSF, in conjunction with dredge surveys. However, adoption of video surveys will require review of the criteria for closures in the fishery to account for the difference in selectivity of dredge surveys compared to video surveys. Dredge surveys, both industry standard and modified to account for recruits, could be conducted in parallel with video surveys for a pre-determined period to correctly calibrate closure criteria of the latter. It is also recommended that concurrent with this use of the technique, further development of the video survey method is undertaken, such as methods to reduce the processing costs of video analysis through techniques such as machine learning. These recommendations will be undertaken in the recently funded FRDC project 2020-030 "Wider investigation of the use of video survey techniques to determine commercial scallop abundance in inshore and offshore waters, closed areas and juvenile beds" commenced in January 2022.

9. References

- Beukers-Stewart, B.D., Vause, B.J., Mosley, M.W.J., Rossetti, H.L., Brand, A.R., 2005. Benefits of closed area protection for a population of scallops. *Mar. Ecol. Prog. Ser.* 298, 189-204.
- Bradshaw, C., Veale, L.O., Hill, A.S., Brand, A.R., 2001. The effect of scallop dredging on Irish Sea benthos: experiments using a closed area. *Hydrobiologia* 465, 129-138.
- Caton, A., McLoughlin, K., 2004. Fisheries Status Reports: Status of fish stocks managed by the Australian Government. Bureau of Rural Sciences.
- Currie, D.R., Parry, G.D., 1996. Effects of scallop dredging on a soft sediment community: a large-scale experimental study. *Mar. Ecol. Prog. Ser.* 134, 131-150.
- DPIPWE, 2005. Assessing the ecological sustainability of the Tasmanian scallop fishery. Department of Primary Industries, Water, and Environment, Hobart, p. 50.
- Ewing, G.P., Keane, J.P., Semmens, J.M., 2018. Industry-independent video survey of commercial scallop (*Pecten fumatus*) densities in Great Oyster Bay – May 2017 survey. Institute for Marine and Antarctic Studies, University of Tasmania, Hobart.
- Ewing, G.P., Moreno, D., Kilpatrick, R., Dudgeon, A., Semmens, J.M., 2021. 2021 Tasmanian scallop fishery preseason survey report. Institute for Marine and Antarctic Studies, University of Tasmania, Hobart.
- Ewing, G.P., Semmens, J.M., 2020. 2020 Tasmanian Scallop Fishery Preseason Survey Report. Institute for Marine and Antarctic Studies, University of Tasmania, Hobart.
- Ewing, G.P., Semmens, J.M., Keane, J.P., Parkinson, J., 2016. Tasmanian Scallop Fishery Assessment 2015. Institute for Marine and Antarctic Studies, University of Tasmania, Hobart.
- Ewing, G.P., Semmens, J.M., Parkinson, J., 2017. Tasmanian Scallop Fishery Assessment 2016. Institute for Marine and Antarctic Studies, University of Tasmania, Hobart.
- Haddon, M., Harrington, J.J., Semmens, J.M., 2006. Juvenile scallop discard rates and bed dynamics: testing the management rules for scallops in Bass Strait, Final report to the FRDC on project 2003/017. Fisheries Research and Development Corporation and Tasmanian Aquaculture and Fisheries Institute.
- Hammond, L.S., Black, K.P., Jenkins, G.P., Hodgkinson, R.S., 1994. The influence of water circulation patterns in Bass Strait on recruitment success and stock differences in scallops, Final report to the FRDC on project 87/117. Victorian Institute of Marine Sciences.
- Harrington, J.J., Haddon, M., Semmens, J.M., 2008. Facilitating industry self-management for spatially managed stocks: a scallop case study, Final report to the FRDC on project 2005/027. Institute for Marine and Antarctic Studies, University of Tasmania.
- Harrington, J.J., Haddon, M., Semmens, J.M., 2008a. Facilitating industry self-management for spatially managed stocks: a scallop case study, Final report to the FRDC on project 2005/027. Institute for Marine and Antarctic Studies, University of Tasmania.
- Harrington, J.J., Semmens, J.M., Haddon, M., 2008b. 2008 Commonwealth Bass Strait Central Zone Scallop Fishery Survey. Survey Final Report. Tasmanian Aquaculture and Fisheries Institute. University of Tasmania.
- Harrington, J.J., Semmens, J.M., Haddon, M., 2007. Spatial distribution of commercial dredge fishing effort: application to survey design and the spatial management of a patchily distributed benthic bivalve species. *Mar. Freshwater Res.* 58, 8.
- Hortle, M., Cropp, D., 1987. Settlement of the commercial scallop, *Pecten fumatus* (Reeve) 1855, on artificial collectors in eastern Tasmania. *Aquaculture* 66, 79-95.

- Jenkins, S.R., Beukers-Stewart, B.D., Brand, A.R., 2001. Impact of scallop dredging on benthic megafauna: a comparison of damage levels in captured and non-captured organisms. *Marine Ecology-Progress Series* 215, 297-301.
- Kruse, G.H., 2015. Independent peer review of the stock assessment of commercial scallops (*Pecten fumatus*) in Tasmanian waters. School of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Hobart, p. 19.
- Lyle, J.M., Stark, K.E., Ewing, G.P., Tracey, S.R., 2019. 2017-18 Survey of Recreational Fishing in Tasmania. Institute for Marine and Antarctic Studies, University of Tasmania Hobart, p. 123.
- Lyle, J.M., Stark, K.E., Tracey, S.R., 2014. 2012-13 Survey of Recreational Fishing in Tasmania. Institute for Marine and Antarctic Studies, University of Tasmania Hobart, p. 10.
- Martin, R.B., Young, P.C., MvLoughlin, R.J., 1988. Problems with applying yield-per-recruit techniques to the management of the Bass Strait Scallop Fishery. In: Zacharin, W.F., Joll, L.M. (Eds.), *Proceedings of the Australian Scallop Workshop*, Hobart, Australia, pp. 111-121.
- Mendo, T., Lyle, J.M., Moltschaniwskyj, N.A., Tracey, S.R., Semmens, J.M., 2014a. Habitat Characteristics Predicting Distribution and Abundance Patterns of Scallops in D'Entrecasteaux Channel, Tasmania. *PLOS ONE* 9, e85895. 60.
- Mendo, T., Moltschaniwskyj, N.A., Lyle, J.M., Tracey, S.R., Semmens, J.M., 2014b. Role of density in aggregation patterns and synchronization of spawning in the hermaphroditic scallop *Pecten fumatus*. *Marine Biology*, 161 (12) pp. 2857-2868.
- Minchin, D., 2003. Introductions: some biological and ecological characteristics of scallops. *Aquat. Living Resour.* 16, 521-532.
- Semmens, J.M., Day, R., Moreno, D., Ewing, G.P., 2020. Fishery-independent video survey of commercial scallop (*Pecten fumatus*) densities in scallop survey sub-area 6A of the Tasmanian Scallop Fishery. Institute for Marine and Antarctic Studies, University of Tasmania, Hobart.
- Semmens, J., Day, R.D., Moreno, D., Ewing, G., Kilpatrick, R., 2023. 2021 Fishery-independent video survey of commercial scallop (*Pecten fumatus*) densities in Tasmanian Scallop Fishery Area 5. Institute for Marine and Antarctic Studies Report. University of Tasmania, Hobart.
- Semmens, J.M., Ewing, G.P., Keane, J.P., 2018. Tasmanian Scallop Fishery Assessment 2017. Institute for Marine and Antarctic Studies, University of Tasmania, Hobart.
- Semmens, J.M., Mendo, T.C., Jones, N.A.R., Keane, J.P., Leon, R., Ewing, G.P., Hartmann, K., 2019. Determining when and where to fish: Linking scallop spawning, settlement, size and condition to collaborative spatial harvest and industry in-season management strategies, Fisheries Research and Development Corporation Final Report, Project Number 2012/027. Institute for Marine and Antarctic Studies, University of Tasmania.
- Semmens, J.M., Ovenden, J.R., Jones, N.A.R., Mendo, T.C., Macbeth, M., Broderick, D., Filardo, F., Street, R., Tracey, S.R., Buxton, C.D., 2015. Establishing fine-scale industry based spatial management and harvest strategies for the commercial scallop in south east Australia, Final Report, Project Number 2008/022. Institute for Marine and Antarctic Studies, University of Tasmania
- Young, P.C., Martin, R.B., McLoughlin, R.J., West, G.J., 1988. Variability in spatfall and recruitment of commercial scallops (*Pecten fumatus*) in Bass Strait. *Proceedings of the Australian Scallop Workshop*, pp. 615-638.